

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/359158034>

# Review on the impact of elevated CO<sub>2</sub> concentrations on fruit species in the face of climate change

Article · March 2022

CITATIONS

3

READS

520

3 authors:



**Gerhard Fischer**

National University of Colombia

324 PUBLICATIONS 4,564 CITATIONS

[SEE PROFILE](#)



**Luz Marina Melgarejo**

National University of Colombia

210 PUBLICATIONS 1,136 CITATIONS

[SEE PROFILE](#)



**Helber Enrique Balaguera-López**

National University of Colombia

144 PUBLICATIONS 839 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ecofisiología, nutrición mineral y manejo integrado de plagas y enfermedades en aguacate, curuba, gulupa y tomate de árbol orientados hacia su manejo agronómico, como materia prima para el desarrollo de productos de interés comercial [View project](#)



Development of floral structures of feijoa (*Acca sellowiana* (O. Berg) Burret) under tropical and subtropical climate conditions. [View project](#)

# Review on the impact of elevated CO<sub>2</sub> concentrations on fruit species in the face of climate change

Revisión del impacto de concentraciones elevadas de CO<sub>2</sub> sobre frutales en la era del cambio climático

 Gerhard Fischer<sup>1\*</sup>  Luz Marina Melgarejo<sup>2</sup>  
 Helber Enrique Balaguera-López<sup>1</sup>

<sup>1</sup> Universidad Nacional de Colombia, Facultad de Ciencias Agrarias, Bogotá, Colombia.

<sup>2</sup> Universidad Nacional de Colombia, Facultad de Ciencias, Bogotá, Colombia.

\*Corresponding author: Gerhard Fischer.  
Universidad Nacional de Colombia,  
Facultad de Ciencias Agrarias, Cra. 30  
No. 45-03, Edificio 500, Bogotá,  
Colombia. [gerfischer@gmail.com](mailto:gerfischer@gmail.com)

Received: March 04, 2021  
Approved: December 14, 2021  
Published: March 08, 2022

Subject editor: Jorge Cadena Torres,  
Corporación Colombiana de  
Investigación, [AGROSAVIA]

How to cite this article: Fischer, G.,  
Melgarejo, L. M., & Balaguera-López, H.  
E. (2022). Review on the impact of  
elevated CO<sub>2</sub> concentrations on fruit  
species in the face of climate change.  
*Ciencia y Tecnología Agropecuaria*, 23(2),  
e2475.

[https://doi.org/10.21930/rcta.vol23\\_num2\\_art:2475](https://doi.org/10.21930/rcta.vol23_num2_art:2475)

**Abstract:** Since the industrial revolution, anthropogenic activities have increased atmospheric CO<sub>2</sub> concentration—one of the major causes of global warming—with a 600 to 700 ppm prediction by the end of this century. Orchards and vineyards are critical sustainable production systems that can minimize emissions and sequester carbon within the atmosphere. Information from different databases (i.e., ScienceDirect, Scopus, SciELO, Google Academic, and ResearchGate) was assessed for this literature review. Generally, elevated CO<sub>2</sub> (e-CO<sub>2</sub>) positively affected fruit trees, such as increased photosynthesis, efficient use of water, growth, and biomass. Therefore, in many cases, the yield and the quality of fruits also increased. With an e-CO<sub>2</sub> of 600-750 ppm, most C3 plants will grow 30 % faster. A total of 1,000 ppm of CO<sub>2</sub> will be optimal for the photosynthesis of various plant species. Fruit trees typically grown in Colombia, such as citrus, grapevines, strawberry, papaya, and pitaya, would benefit from these positive effects, as e-CO<sub>2</sub> alleviates stress due to drought and waterlogging. However, the increased growth of fruit trees due to e-CO<sub>2</sub> requires more nutrients and water. Thus, selecting genotypes that benefit from e-CO<sub>2</sub> and have high efficiency in using nitrogen and water is very important. Ideally, they must have a high sink strength to avoid the accumulation of carbohydrates in the chloroplast. The authors conclude that there is undoubtedly a “fertilization effect of CO<sub>2</sub>” on fruit species that increases with the advance of climate change. Still, much research is lacking for fruit species compared to many other crops. Hence, future studies are required to measure the direct effects of atmospheric e-CO<sub>2</sub> and its interactions with environmental variables such as rainfall, temperature, soil moisture, and nutrient availability.

**Keywords:** Carbon dioxide, carbon dioxide enrichment, nitrogen, photosynthesis, sink strength, source sink relations, water use efficiency, fruit trees physiology.

**Resumen:** Las actividades antropogénicas han contribuido a que la concentración de CO<sub>2</sub> atmosférico aumente constantemente con una predicción de 600 a 700 ppm para fines de este siglo, siendo una de las mayores causas del calentamiento global. Los huertos frutales y viñedos son importantes sistemas de producción sostenible que pueden minimizar las emisiones y secuestrar carbono de la atmósfera. Para esta revisión de literatura, se evaluó mediante la información obtenida de diferentes bases de datos. Generalmente, el CO<sub>2</sub> elevado (e-CO<sub>2</sub>) genera efectos positivos sobre los frutales en procesos como el aumento de la fotosíntesis, el uso eficiente de agua, el crecimiento y la biomasa. Por lo anterior, en muchos casos, el rendimiento y la calidad de los frutos también incrementaron. Se estima que, con un e-CO<sub>2</sub> de 600-750 ppm, la mayoría de las plantas C3 crecerán un 30 % más rápido. Con 1000 ppm las condiciones serán óptimas para la fotosíntesis de varias especies vegetales. Los árboles frutales que también crecen en Colombia como los cítricos, la vid, la fresa, la papaya y la pitaya, se beneficiarían de los efectos positivos mencionados anteriormente, en tanto que el e-CO<sub>2</sub> aliviaría los efectos del estrés por sequía y anegamiento. Sin embargo, el mayor crecimiento de los frutales por el e-CO<sub>2</sub> exige un mayor suministro de nutrientes y agua, por lo cual es muy importante la selección de genotipos que se beneficien del e-CO<sub>2</sub> y que presenten un alto uso eficiente de nitrógeno y agua. Así mismo, es deseable que dichas especies posean una alta fuerza vertedero para evitar la acumulación de carbohidratos en el cloroplasto. Esta revisión permite concluir que existe un “efecto fertilizante del CO<sub>2</sub>” sobre las especies frutales que aumenta con el avance del cambio climático. Sin embargo, existe poca investigación en comparación con muchos otros cultivos agrícolas. Por ello, a futuro se requieren estudios que midan los efectos directos del e-CO<sub>2</sub> atmosférico y sus interacciones con variables ambientales, como la lluvia, la temperatura, la humedad del suelo y la disponibilidad de nutrientes.

**Palabras clave:** dióxido de carbono, enriquecimiento de dióxido de carbono, fotosíntesis, fuerza vertedero, nitrógeno, uso eficiente del agua, fisiología de frutales, relaciones fuente sumidero.



## Introducción

Uncontrolled deforestation and increasing greenhouse gas emissions are anthropogenic activities responsible for triggering environmental disequilibrium that affects the Earth's complex climate dynamism (Menezes-Silva et al., 2019). The carbon dioxide (CO<sub>2</sub>) concentration in the atmosphere has risen at a rate of  $2.16 \pm 0.09$  ppm year<sup>-1</sup> in recent decades as a result of human activity (Le Quéré et al., 2018), and its increase is expected to continue until emissions are reduced (Anderson et al., 2019). Kumar et al. (2017) reported that CO<sub>2</sub> levels had reached concentrations above 400 μmol/mol, and a record increase of 415 ppm has been reported recently (Brito et al., 2020). By the end of this century, the prediction of 600 to 700 ppm will increase the mean surface temperature from 4.5 to 5.0 °C (Leung et al., 2014) and CO<sub>2</sub> concentration by 1 to 3 ppm per year (Taiz et al., 2017). The increase in global temperature has a direct impact on the structure and function of ecosystems, including its effects on the physiology and growth of trees (Ceulemans et al., 1999), with CO<sub>2</sub> being the critical substrate for photosynthesis and, therefore, the main contributor to world food production (Mishra et al., 2019).


Of the CO<sub>2</sub> entering the atmosphere each year due to human activity, 45 % remains there, while oceans sequester 25 %, and terrestrial ecosystems the rest at 30 % on average (Henson, 2011). However, there are significant variations from one year to the next due to climatic cycles and changes in land use (Henson, 2011).

Apart from CO<sub>2</sub>, emissions associated with greenhouse gases (GHG) mainly include methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). The three add up to 80 % of the GHG (IPCC, 2013). Furthermore, GHGs are associated with fluorocarbons (HFCs), perfluorocarbons, and sulfur hexafluoride (SF<sub>6</sub>) (López-Bellido, 2015). These GHGs have increased dramatically in recent decades (Al-Mamoori et al., 2017; Ouda et al., 2016) caused mainly by the combustion of fossil materials (coal, oil, and natural gas), industrial emissions, deforestation, and soil degradation (Henson, 2011; IPCC, 2013).

According to the IPCC (2019), climate change affects food security due to warming, altered rain patterns, and a greater frequency of extreme weather events. Furthermore, the impact and consequences of climate change for agriculture tend to be more severe for countries with higher initial temperatures, areas with marginal or already degraded lands, and lower levels of development with little adaptive capacity (Yohannes, 2016). Mishra et al. (2019) stated that in the long term, the exposure of plants to high CO<sub>2</sub> (e-CO<sub>2</sub>), high temperatures, and droughts would considerably affect the balance of ecosystem processes at the local and global levels.

The e-CO<sub>2</sub> in trees markedly improves productivity due to increased efficiency in water use, a high rate of photosynthesis, more sugar accumulation in fruits, and higher biomass production (Rajan et al., 2020) (figure 1). However, Ebi et al. (2021) lessen this enthusiasm because e-CO<sub>2</sub> can also alter the nutritional quality of C3 plants. Fruit plantations can contribute positively to sustainable development in the climate change scenario in the tropics, considering that the current expansion of agriculture and poverty level continues reducing forest resources (Patil & Kumar, 2017). Wu et al. (2012) estimate that apple orchards in China, the largest producer of

this fruit globally, can offset between 1.6 % and 3.0 % of CO<sub>2</sub> emissions from burning fossil fuels in China, highlighting the importance of these orchards for carbon sequestration.



**Figure 1.** Diagram of the increase (↑) and decrease (↓) of growth parameters and physiology of the fruit plant under increased atmospheric CO<sub>2</sub>.

Source: Modified based on Mishra et al. (2019) and Ramírez and Kallarackal (2015)

There is enough research that indicates that atmospheric e-CO<sub>2</sub> increases net photosynthesis, biomass accumulation, seed and fruit yield, water use efficiency (figure 1), light interception, nutrient uptake, and the water potential of plants (Tognetti et al., 2005). Overall, e-CO<sub>2</sub> is expected to improve the fruit nutrient content (Balasooriya et al., 2019). Nevertheless, the studies that find interactions with other environmental and genetic factors that modify these results in e-CO<sub>2</sub> events should not be underestimated (Mishra et al., 2019).

Thus, plant responses to e-CO<sub>2</sub> and the resulting changes in species can be strongly affected by interactions with other environmental factors and climate change, including tropospheric ozone, temperature, water, and nitrogen (Bradley & Pregitzer, 2007). For this reason, Allen and Vu (2009) warned that global e-CO<sub>2</sub> does not have the same effects on all plant species and all environments, so it is unwise to extend predictions from research in well-irrigated systems to those cultivated in hot and arid environments or milder climatic zones.

In many cases, the effects of climate change are examined through each factor involved, but it must be considered that the response of plants to climate change is not the sum of the reactions to the different phenomena but results from the interaction between all these (Pérez-Jiménez et al., 2017). According to the concept of multidimensionality by Zandalinas et al. (2021), all climatic factors occur at the same time, which is why studies in open areas such as the FACE system (Free Air CO<sub>2</sub> Enrichment) (Ramírez & Kallarackal, 2015; Sánchez et al., 2015) or also semi-open (open-top chambers) have gained much importance in research into climate change in our crops. Studies of an e-CO<sub>2</sub> in the FACE system, which usually runs for several years, have shown effects on photosynthesis and stomata, kept closed longer to directly

reduce transpiration (figure 1). However, mitochondrial respiration increases due to increased leaf temperature (Nobel, 1999; Taiz et al., 2017).

Nobel (1991) stated that many C<sub>3</sub> plants (all fruit trees except pineapple and cacti such as pitaya and prickly pear, which are CAM plants) need an optimal CO<sub>2</sub> concentration of about 1,000 ppm to fix carbon during photosynthesis since no saturation has been observed at the current levels of ambient CO<sub>2</sub> (Mishra et al., 2019). Taiz et al. (2017) reported that with e-CO<sub>2</sub> up to 600-750 ppm, most C<sub>3</sub> plants would grow 30 % faster, but the growth rate may become limited by the availability of nutrients for the plant (figure 1).

Because the current concentration of CO<sub>2</sub> barely exceeds 400 ppm (Kumar et al., 2017), CO<sub>2</sub> deficiency is one of the limiting factors for photosynthesis and crop production (Fischer et al., 2016; Song et al., 2020). An atmospheric e-CO<sub>2</sub> would increase plant photosynthesis through a “CO<sub>2</sub> fertilizer effect” (Mishra et al., 2019). This effect is generated by biomass production (figure 1) derived from photosynthesis, defined by Larcher (2003) as 85 % to 92 % of dry matter. Due to the more significant accumulation of carbohydrates by plants, e-CO<sub>2</sub> will also increase its availability for symbiotic organisms such as mycorrhizal fungi and associated rhizospheric bacteria (Bhargava & Mitra, 2021).

The e-CO<sub>2</sub> up to 1,000 μmol/mol used inside greenhouses to increase crop production (mostly of vegetables and strawberries) is a technology that has been applied for several years (Becker & Kläring, 2016). However, in greenhouses, especially for reasons of the indoor climate, ventilation cannot be opened. Therefore, the concentrations of CO<sub>2</sub> in the air can decrease to 150 μmol/mol during the day due to plants, as found by Kläring et al. (2007) for cucumber.

Nitrogen availability and cycling play a crucial role in regulating responses to e-CO<sub>2</sub>, especially in temperate ecosystems (Reich et al., 2006) but also in the tropics (DaMatta et al., 2010) because low N levels do not stimulate increased production in systems with CO<sub>2</sub> enrichment (figure 1) (Leakey et al., 2012). Thus, they alter the relationship between nutrient demand and crop growth (Cruz et al., 2016). In this context, it is essential to consider that more significant applications of nitrogen fertilizers can increase the release of N<sub>2</sub>O into the atmosphere, which is a potent greenhouse gas. Therefore, efforts should increase efficiency in using nitrogen from plants (Jackson et al., 2011). It is also true that moderate CO<sub>2</sub> enrichment can improve N absorption efficiency and decrease N loss from the soil associated with a decrease in nitrification and denitrification under a high application of N (Dong et al., 2020).

Experiments in temperate zones showed that e-CO<sub>2</sub> could alleviate drought stress (Leakey, 2009) and the drying effects of warming (Morgan et al., 2011). In general, it is assumed that water use in the plant canopy is lower under e-CO<sub>2</sub> conditions because, although the leaf area index may increase, there is a lower stomatal conductance on average (Ainsworth & Lemonnier, 2018) that compensates by a larger evaporation surface (Ainsworth & Rogers, 2007). In tomato, Wei et al. (2018) found that an e-CO<sub>2</sub> of 800 ppm attenuated the effects of water-deficit stress on performance.

It is well known that high temperatures reduce the net gain of carbon in C<sub>3</sub> species by increasing photorespiration (Nobel, 1999). Therefore, e-CO<sub>2</sub> can decrease photorespiration

and increase photosynthesis, mainly under high-temperature conditions rather than low temperatures (DaMatta et al., 2010), partially offsetting the effects of supra-optimal temperatures on yield (Polley, 2002). DaMatta et al. (2010) concluded that e-CO<sub>2</sub> would have the most positive effects on the growth of crops with temperatures close to the optimum. In regions highly affected by high temperatures, such as low latitudes, a progressive increase in heat decreases crop yield independently of e-CO<sub>2</sub> (Polley, 2002).

Usually, the e-CO<sub>2</sub> applied by horticulturists generates higher yields (Gruda et al., 2019). However, ambient CO<sub>2</sub> can reduce input costs for CO<sub>2</sub> enrichment. Gruda et al. (2019) reported it as unlikely that these applications will be unnecessary in the future. Crops benefit from even higher levels with this gas, considering the most pessimistic scenario of the IPCC (2013) for 2100 with about 935 ppm of CO<sub>2</sub>. On the other hand, CO<sub>2</sub> applications in greenhouses are restricted by the ambient temperature. If it is high, the vents must be opened, making the e-CO<sub>2</sub> ineffective due to losses to the outside (De Zwart, 2012). This effect may still be aggravated by heat waves generated by climate change (Bisbis et al., 2018).

There are few publications on fruit trees with e-CO<sub>2</sub> than other crops (Ramírez & Kallarackal, 2015; Wohlfahrt et al., 2018). Studies, especially on vegetables, showed that with e-CO<sub>2</sub>, the concentrations of calcium, glucose, fructose, total soluble solids, total flavonoids, total phenols, ascorbic acid, and the total antioxidant capacity could increase in the edible part of vegetables. However, e-CO<sub>2</sub> can also decrease the concentrations of nitrate, magnesium, zinc, iron, and protein (Dong et al., 2018).

Becker and Kläring (2016) found that e-CO<sub>2</sub> in greenhouses or plant factories for species such as lettuce increases the availability of precursors for the biosynthesis of phenolic compounds that are health promoters. Increased e-CO<sub>2</sub> can also increase the nutritional quality of fruits (Fischer et al., 2016). Moretti et al. (2010) reported that e-CO<sub>2</sub> increased ascorbic acid levels in postharvest strawberries and oranges. Therefore, this review article aims to report the effects of the increase in atmospheric CO<sub>2</sub> on the physiology, growth, and production of fruit species. This knowledge can be used to decide on species selection, management, and the mechanisms to mitigate these effects.

## Materials and methods

The information from different databases, including ScienceDirect, Scopus, SciELO, Google Academic, and ResearchGate, was assessed for this literature review. The search was made using keywords (in English and Spanish) such as “CO<sub>2</sub>,” “carbon dioxide,” “CO<sub>2</sub> fertilization,” “carbon fertilization,” “climate change,” “fruits,” “fruit trees,” “fruit plants.” From the mentioned databases, we obtained 88 sources that include scientific national and international articles, books, and book chapters from the last 30 years, in English and Spanish.

The different fruit trees were chosen according to the research with CO<sub>2</sub> in the last three decades. From these, the studies of species important in Colombia, such as different citrus fruits, grape, papaya, strawberry, and pitaya, are described in more detail in table 1.

## Effect of high CO<sub>2</sub> on the physiology and production of fruit trees

The effects of e-CO<sub>2</sub> in the air near the leaf blade are well-known and reported. They include decreased stomatal opening, stomatal conductance, and transpiration. In response, photosynthesis and plant growth increase. Additionally, the plants show greater efficiency in the use of water (Pritchard & Amthor, 2005; Stöckle et al., 2011) and light (Drake & González-Meler, 1997). The key reason for this enriched photosynthesis is the increased carboxylation efficiency of RuBisCO, which is relatively low in the concentration of ambient atmospheric CO<sub>2</sub> (Mishra et al., 2019).

If the environmental level of CO<sub>2</sub> increases from 350 to 550 ppm (at 25 °C), with time, the rates of photosynthesis will be reduced in some species, compared to plants grown at ambient levels of CO<sub>2</sub>. This effect is called “photosynthetic acclimation,” attributed to five mechanisms at the cellular level, reported by Ramírez and Kallarackal (2015): (1) gene repression and sugar accumulation; (2) insufficient nitrogen uptake by the plant; (3) a link of carbohydrate accumulation with inorganic phosphate and, consequently, a limitation in the renewing capacity of RuBP; (4) accumulation of starch in the chloroplast; and (5) capability of triose phosphate utilization. Photosynthetic acclimation with e-CO<sub>2</sub> is a critical topic that must be assessed in detail because it would indicate that the advantage of e-CO<sub>2</sub> can be lost over time. Therefore, new studies must involve several growing cycles.

Since e-CO<sub>2</sub> generates higher photosynthesis rates, a more significant amount of assimilates is available to be channeled towards biosynthetic pathways of different types (Treutter, 2010), resulting in higher levels of secondary metabolites due to the greater availability of precursor molecules (Becker & Kläring, 2016). Furthermore, in e-CO<sub>2</sub> environments, because of the greater number of growing sink organs, there is an increase in the demand for photo-assimilates, which stimulate the net assimilation rates of carbon in the form of sucrose and starch (Bhargava & Mitra, 2021). e-CO<sub>2</sub> (750 ± 30 μmol/mol) can increase photosynthesis rates without affecting stomatal conductance and dark respiration. Nevertheless, ABA and 1-aminocyclopropane-1-carboxylic acid levels cannot do this without a stressful condition such as salinity (reported in the tomato), where the intermediates of the Krebs cycle in e-CO<sub>2</sub> can also be increased to reduce the harmful effect of salinity (Brito et al., 2020). However, e-CO<sub>2</sub> can also decrease the respiratory rate (Kochhar & Gujral, 2020). Cytochrome pathway respiration may be affected, and respiration may be increased through the alternative oxidase pathway due to a decrease in O<sub>2</sub>, an aspect that also deserves more attention.

Bitter orange trees planted in open-top chambers with transparent plastic for 17 years at 300 ppm CO<sub>2</sub> higher than the ambient concentration (table 1) showed a 70 % increase in the total biomass of the orange tree compared to the control tree (Kimball et al., 2007). This improvement came from the more significant number of fruits produced, mainly by numerous branches, an increase in the thickness of the trunk and branches, and an increase in the length of the fine roots in 64.5 % and 57.2 % at soil depths of 0-15 and 15-30 cm, respectively (Prior et al., 2012). These advantages can considerably improve the productivity of fruit trees.

**Table 1.** Examples of the effect of CO<sub>2</sub> enrichment on fruit trees

Crop	Treatment	Response	Author
Orange tree ‘Ambersweet’ (‘Clementine’ tangerine × ‘Orlando’ tangelo)	Greenhouses with 360 and 720 ppm CO <sub>2</sub> , at 0, 1.5, 3.0, and 4.5 °C above ambient temperature.	The e-CO <sub>2</sub> increased the growth parameters except for leaves and fine root biomass. There was no CO <sub>2</sub> × temperature interaction.	Allen and Vu (2009)
Bitter orange ( <i>Citrus aurantium</i> )	Plants in open-top chambers with transparent plastic, for 17 years at 300 ppm CO <sub>2</sub> higher than the ambient concentration.	The e-CO <sub>2</sub> increased the total biomass of the orange tree by 70 %, noted in a more significant number of fruits and branches. It increased the thickness of the trunk and branches and the number of roots. There were no significant changes in the elemental composition of the biomass produced.	Kimball et al. (2007)
Satsuma mandarin ( <i>Citrus unshiu</i> cv. Uenowase)	Photosynthesis of the fruit bark and leaves at concentrations between 50 and 1,000 ppm CO <sub>2</sub> was measured.	Up to 500 ppm e-CO <sub>2</sub> , the gross photosynthetic rate of the fruit shell increased, but at higher concentrations, it decreased. Foliar photosynthesis constantly increased between 50 and 1,000 ppm CO <sub>2</sub> .	Hiratsuka et al. (2015)
Lemon ( <i>Citrus limon</i> var. ‘Villafranca,’ grafted onto <i>C. volkameriana</i> )	Three greenhouse compartments at 400, 650, and 850 ppm CO <sub>2</sub> , with water-deficit stress of the plants for one month (after two months of irrigation), followed by re-irrigation for another month.	Seedlings exposed to e-CO <sub>2</sub> and a 1-month drought maintained their growth but decreased at 400 ppm. The down-regulation of stomata to e-CO <sub>2</sub> decreased plants’ foliar transpiration and daytime water use by 13 % - 46 %, but with even higher photosynthesis by 15 % - 25 % than those grown at ambient CO <sub>2</sub> .	Paudel et al. (2018)
( <i>Prunus cerasus</i> × <i>P. canescens</i> ) sweet cherry rootstock	400 and 800 ppm CO <sub>2</sub> , 26 °C, with and without waterlogging.	e-CO <sub>2</sub> increased photosynthesis and non-structural carbohydrates, helping plants overcome waterlogging.	Pérez-Jiménez et al. (2017)
Grape ( <i>Vitis vinifera</i> ) ‘Tempranillo’	In greenhouses with ambient CO <sub>2</sub> at 24/14 °C and 700 ppm CO <sub>2</sub> at 28/18 °C (climate change), UV-B 0, 5.98, and 9.66 kJ m <sup>-2</sup> d <sup>-1</sup> .	Climate change always generated the highest photosynthetic yield. UV-B can help alleviate the signs of oxidative damage and slow berry ripening under high temperature and CO <sub>2</sub> conditions.	Martínez-Lüscher et al. (2015)
Grape ( <i>Vitis vinifera</i> ) ‘Tempranillo’ red and white	In a greenhouse with two levels of CO <sub>2</sub> (400 and 700 ppm) × two temperatures (ambient and 4 °C higher) × two water regimes (well-watered and cyclical drought).	The yield was significantly reduced by drought. Thermal shocks (> 35 °C) induced burns and losses of 50 % of the berries. High temperatures and drought increased the pH of the grape juice due to the decrease in malic acid; e-CO <sub>2</sub> decreased the pH with increases in tartaric acid.	Kizildeniz et al. (2018)





Grape ( <i>Vitis vinifera</i> ) 'Riesling' and 'Cabernet Sauvignon'	FACE with ambient CO <sub>2</sub> (400 ppm) and e-CO <sub>2</sub> (+ 20 % of ambient CO <sub>2</sub> ).	Increase in primary productivity in e-CO <sub>2</sub> due to greater assimilation of CO <sub>2</sub> . WUE <sub>i</sub> , transpiration rate, stomatal conductance, and pre-dawn leaf water potential increased under e-CO <sub>2</sub> . The e-CO <sub>2</sub> resulted in a higher bunch weight without altering the sugar content of the juice on the date of harvest.	Wohlfahrt et al. (2018)
Strawberry ( <i>Fragaria ananassa</i> cv. Toyonoka)	At two temperatures, 20/15 °C, and 25/20 °C day/night at 360 and 720 ppm CO <sub>2</sub> with 0 and 0.01 % NH <sub>4</sub> NO <sub>3</sub> in growth chambers.	The e-CO <sub>2</sub> and high temperature caused a further decrease of 12 % and 35 % in fruit yield, respectively, regardless of fertilization with low or high nitrogen concentrations. The e-CO <sub>2</sub> increased dry matter, fructose, glucose, total sugar, and sweetness index but decreased the total antioxidant capacity in fruits. Low-temperature e-CO <sub>2</sub> increased the production and quality of the strawberry.	Sun et al. (2012)
Strawberry ( <i>Fragaria ananassa</i> var. 'Hongyan')	e-CO <sub>2</sub> (750-850 ppm) <i>vs.</i> environmental CO <sub>2</sub> (350-500 ppm) in the Chinese Solar Greenhouse compartments.	The e-CO <sub>2</sub> increased the size, net photosynthetic rate, and light saturation point of the leaves but decreased the light compensation point. The foliar tissue transcriptome identified 150 genes differentially expressed in response to e-CO <sub>2</sub> , of which 14 genes were involved in photosynthesis.	Li et al. (2020)
Papaya ( <i>Carica papaya</i> ) Tainung #1 F1 híbrido	e-CO <sub>2</sub> of 750 ppm <i>vs.</i> control (390 ppm) in the greenhouse and two levels of nitrogen (3 mM or 8 mM).	The e-CO <sub>2</sub> increased the assimilation rate of the seedlings at both levels of N. The e-CO <sub>2</sub> increased the instantaneous efficiency of water use with the two levels of N. The e-CO <sub>2</sub> stimulated the accumulation of dry mass in plants, especially for plants cultivated with lower levels of N.	Cruz et al. (2016)
Yellow pitaya ( <i>Selenicereus megalanthus</i> )	e-CO <sub>2</sub> of 1,000 ppm <i>vs.</i> control (380 ppm) in heated greenhouse chambers.	Plants with e-CO <sub>2</sub> of 1,000 ppm reacted with 129, 68, and 233 % increases in the daily intake of CO <sub>2</sub> , dry mass of the stem, and the number of flower buds, respectively. In addition, fruits' fresh weight increased by 63 %, compared to those grown in environmental CO <sub>2</sub> .	Weiss et al. (2010)

Source: Elaborated by the authors.

Young lemon (*Citrus limon*) plants exposed to 650 and 850 ppm CO<sub>2</sub> and a 1-month drought maintained their growth. However, at 400 ppm it decreased (table 1) due to the down-regulation of the stomata by which the foliar transpiration and the daytime use of water of the plants decreased by 13 % - 46 %, although the photosynthesis was 15 % - 25 % higher compared to environmental CO<sub>2</sub> (Paudel et al., 2018). The authors concluded that e-CO<sub>2</sub> partially offset the effects of drought on plant development and thus diminished some of the effects of anthropogenic climate change.

In the Satsuma mandarin with e-CO<sub>2</sub>, up to 500 ppm increases the gross photosynthetic rate of the fruit peel, but at higher concentrations, the rate decreases, a reaction similar to C<sub>4</sub> photosynthesis (Hiratsuka et al., 2015). The authors find that foliar photosynthesis constantly increases between 50 and 1,000 ppm CO<sub>2</sub> (table 1). Interestingly, the photosynthesis of the fruit was higher than the foliar at a low photosynthetic photon flux density (PPFD) of 13.5 to 68  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ; for this reason, Hiratsuka et al. (2015) characterized the species as an intermediate state between C<sub>3</sub>, C<sub>4</sub>, and shade plants.

Viticulture regions are confined to unique climatic niches; climate change affects them significantly due to changes in precipitation and temperature. The changes can destabilize the balance between climate, soil, and plant, with profound effects on the production of high-quality wines (Moriondo et al., 2013), meaning global warming will shift for new regions towards the poles and in the tropics to the increased altitude that may potentially become more suitable for growing and producing high-quality wines (Jones et al., 2005). This situation is critical in Colombia, where grapevines are grown above 2,200 m a.s.l.

The greenhouse and FACE studies (table 1) showed an increase in photosynthetic yield and thus a higher production of grapes (Martínez-Lüscher et al., 2015; Wohlfahrt et al., 2018). Bindi et al. (2001) observed a 45–50 % increase in grape biomass production when the CO<sub>2</sub> concentration rose to 700  $\mu\text{mol/mol}$  without adverse effects on the quality of the grape or wine. However, when e-CO<sub>2</sub> was accompanied by drought or thermal shocks due to supra-optimal temperatures, induced burns and a considerable loss of the berries greatly affected the yield (table 1) (Kizildeniz et al., 2018).

Performance is highly dependent on photosynthesis, and CO<sub>2</sub> is one of the essential constituents of this process (Li et al., 2020). In strawberries, e-CO<sub>2</sub> of 720 ppm and high temperatures (25/20 °C day/night) decreased the yield due to the lower number of inflorescences induced at these temperatures, while low temperatures (20/15 °C) increased their production and quality (table 1) (Sun et al., 2012). These authors warned that the optimal temperatures for flower induction must be considered for e-CO<sub>2</sub>, which must be low since global warming can affect strawberry production.

Effects on the quality of strawberries occur when e-CO<sub>2</sub> (300 and 600 ppm higher than ambient CO<sub>2</sub>) increases the concentrations of flavonoids and anthocyanins in fruits (Wang et al., 2003). The authors found that plants grown in e-CO<sub>2</sub> showed higher oxygen-free radical absorbance activity in the fruit (Wang et al., 2003). Two studies of the effect of e-CO<sub>2</sub> on strawberry leaves showed an increase in leaf size, photosynthesis, and light saturation (table 1) (Li et al., 2020). These results confirmed what was observed by Keutgen et al. (1997) in which

an e-CO<sub>2</sub> of 600 ppm promoted net foliar photosynthesis, while at a higher concentration, it led to a decrease in net assimilation.

Cactaceous fruits such as yellow pitaya (*Selenicereus megalanthus*), characterized by the acid metabolism of Crassulaceae (CAM), reacted to an e-CO<sub>2</sub> of 1,000 ppm (*vs.* 380 ppm environmental) with a tremendous increase in the daily intake of CO<sub>2</sub> and vegetative and reproductive growth factors (table 1) compared to the red pitaya (*Hylocereus undatus*) that also showed a considerable increase in vegetative growth, but only a 7 % increase in the fresh weight of fruits, compared to 63 % of yellow pitaya (Weiss et al., 2010). The authors attributed the response to a differential adaptation of these two species to humid conditions (Weiss et al., 2010). Rajan et al. (2020) classified pitayas (dragon fruits) as very suitable for climate change conditions due to their lower demand for soil moisture and reduced transpiration rate (figure 1).

CO<sub>2</sub> fixation occurs mainly at night in CAM plants, catalyzed by the cytosolic enzyme phosphoenolpyruvate carboxylase (PEPC) to produce malate or aspartate, stored in vacuoles (Mishra et al., 2019; Nobel, 1999). Decarboxylation occurs during the day and results in the release of CO<sub>2</sub> from malic or aspartic acid and its conversion to carbohydrates in the Calvin cycle (Taiz et al., 2017). Due to their transpiration efficiency (three to five times higher than C3 or C4 plants), these plants are more suitable in places with water scarcity (Mishra et al., 2019).

Cruz et al. (2016) found that for papaya seedlings, e-CO<sub>2</sub> at 750 ppm increased not only the assimilation rate but also the instantaneous efficiency of water use (table 1), especially at the higher level of N applied (52 % at 8 mM N), compared to the lowest (16 % at 3 mM N). The e-CO<sub>2</sub> stimulated the accumulation of dry mass in the plants, especially for plants cultivated with lower levels of N. However, e-CO<sub>2</sub> decreased N concentrations in all plant organs (figure 1), regardless of the level of N used (Cruz et al., 2016).

Unfortunately, neither Colombia nor other countries have studied CO<sub>2</sub> on important fruits for export, *e.g.*, cape gooseberry, purple passionfruit, and sweet granadilla. In Colombia, no studies have been carried out with different concentrations of CO<sub>2</sub> in fruit trees.

### **Aspects of mitigating the adverse effect of CO<sub>2</sub> on fruit trees**

Despite the increase in atmospheric CO<sub>2</sub>, considering global warming and altered rainfall, food production is uncertain for the future (Haokip et al., 2020). For the profitable production of fruit trees, climate change, especially e-CO<sub>2</sub>, is a tremendous challenge for the producer needing to find possible solutions to mediate these effects (Fischer et al., 2016). Therefore, fruit growers must achieve high efficiency in the use of nutrients (particularly nitrogen) and water, while guaranteeing a sufficient entry of light (Ramírez & Kallarackal, 2015), *e.g.*, through greater distances between trees, so that plant conduction and pruning achieve optimal leaf growth and development (Casierra-Posada & Fischer, 2012).

Fruit growers must guarantee adequate “soil fertilization with CO<sub>2</sub>” to increase soil respiration (Fischer et al., 2016). In Italy, organically managed vineyards that apply manure and bury pruning residues showed higher soil respiration rates than conventional orchards (Brunori et

al., 2016). Thus, a 2/3 increase in soil CO<sub>2</sub> production is seen for microorganisms and 1/3 for root respiration when fertilizers are applied organically (Fischer & Orduz-Rodríguez, 2012).

Likewise, producers must mitigate climate change by selecting more adapted cultivars that respond to e-CO<sub>2</sub> and are capable of tolerating drought or waterlogging conditions (Fischer et al., 2016). The plant's response capacity to e-CO<sub>2</sub> is related to the sink's intrinsic capacity, adaptive plasticity, and good progress. These aspects could be achieved with plant breeding programs or genetic engineering (Dingkuhn et al., 2020). Also, Mishra et al. (2019), with a view to prolonged e-CO<sub>2</sub> levels (the reality of climate change), underline the species and varieties with high sink strength to accumulate carbohydrates. These plants do not react with suppression of photosynthesis, unlike the plants that accumulate carbohydrates excessively in leaf tissues (figure 1). In this regard, exploring the genomic basis of local adaptation is essential for evaluating the conditions under which fruit trees will successfully adapt *in situ* to global climate change (Cortés et al., 2020).

Because e-CO<sub>2</sub> is accompanied by an increase in temperature and, in many cases, by more intense dry seasons, new orchards should be established at higher altitudes and latitudes (Fischer & Melgarejo, 2020; Houston et al., 2018) where there is sufficient availability of water for irrigation. Preferably irrigation systems should be used, which reduce GHG emissions (CO<sub>2</sub> and N<sub>2</sub>O), such as underground drip irrigation (Jackson et al., 2011) or deficit irrigation that saves much water without damaging the quality of fruits (Vélez-Sánchez et al., 2019, 2021). An increase in altitude decreases CO<sub>2</sub> partial pressure and H<sub>2</sub>O vapor (Nobel, 1999). However, in the case of deciduous fruit trees in the inner tropics, it is not sure that an increase in altitude for the plantations will guarantee sufficient chilling hours to break bud rest so that the growers continue defoliating and applying chemicals to trees to break dormancy (Luedeling et al., 2011).

Based on observations of farmers' performance in the face of extreme climatic events, resilience is highly related to the high biodiversity of farms, typical in traditional agricultural systems, especially in combination with agroecological methods that are the only viable methods for maintaining productivity and sustainability of peasant agriculture under climate change (Altieri & Nicholls, 2017; Pérez et al., 2010). However, further studies are required under the new scenarios to evaluate the impacts of climate change, as they may serve as tools in decision-making for the management of fruit species and their productivity.

To avoid losses in fruit production due to e-CO<sub>2</sub>, Haokip et al. (2020) proposed implementing a plan based on a strategic scientific impact assessment with the adaptations and mitigation necessary. Sharma et al. (2021) confirmed that sustainable fruit production systems could mitigate emissions and sequester carbon within the atmosphere. Besides the shallow soil tillage, which preserves the soil organic matter, the structural characteristics of orchards and vineyards, including their long-life cycle, permanent organs (trunk, branches, roots), and high yields, force them to accumulate a considerable amount of carbon (Sharma et al., 2021). Also, Marín et al. (2016) reported high carbon fixation rates (17.7 t year<sup>-1</sup>) in agroforestry systems with citrus species.

## Conclusions

Due to human activity, the concentration of CO<sub>2</sub> is constantly increasing and stands out as one of the major causes of global warming. Generally, e-CO<sub>2</sub> positively affects fruit trees, such as increased photosynthesis, water use efficiency, and growth. Therefore, in many cases, the yield and the quality of fruits also increase.

The increased growth of fruit trees due to e-CO<sub>2</sub> requires more nutrients and water, so selecting genotypes that benefit from e-CO<sub>2</sub>, make highly efficient use of nitrogen and water, and have a high sink strength, is very important. More importantly, fruit growers should select cultivars adapted to climate change that respond to e-CO<sub>2</sub> and tolerate drought or waterlogging conditions.

Thanks to their permanent organs, sustainable fruit production systems can minimize emissions and sequester carbon within the atmosphere. Also, to fulfill the requirement of sufficient CO<sub>2</sub> fertilization, especially of the lower plant part, increased production of soil CO<sub>2</sub> through organic fertilizer application is a good strategy.

The authors conclude that there is unquestionably a “fertilization effect of CO<sub>2</sub>” on fruit species that increases with the progress of climate change. However, much research lacks fruit trees compared to many other field crops. Further studies are required to address the direct effects of atmospheric e-CO<sub>2</sub> and its interactions with environmental variables such as rainfall, temperature, soil moisture, nutrient availability, and the vapor pressure deficit with altitude.

## Disclaimers

All the authors made significant contributions to the document, agree with its publication, and state no conflicts of interest in this study.

## References

- Ainsworth, E. A., & Lemonnier, P. (2018). Phloem function: a key to understanding and manipulating plant responses to rising atmospheric [CO<sub>2</sub>]? *Current Opinion in Plant Biology*, 43, 50-56. <https://doi.org/10.1016/j.pbi.2017.12.003>
- Ainsworth, E. A., & Rogers, A. (2007). The response of photosynthesis and stomatal conductance to rising [CO<sub>2</sub>]: mechanisms and environmental interactions. *Plant Cell and Environment*, 30, 258-270. <https://doi.org/10.1111/j.1365-3040.2007.01641.x>
- Al-Mamoori, A., Krishnamurthy, A., Rownaghi, A. A., & Rezaei, F. (2017). Carbon capture and utilization update. *Energy Technology*, 5(6), 834-849. <https://doi.org/10.1002/ente.201600747>

- Allen, L. H., & Vu, J. C. V. (2009). Carbon dioxide and high temperature effects on growth of young orange trees in a humid, subtropical environment. *Agriculture and Forest Meteorology*, 149, 820-830. <https://doi.org/10.1016/j.agrformet.2008.11.002>
- Altieri, M. A., & Nicholls, C. I. (2017). The adaptation and mitigation potential of traditional agriculture in a changing climate. *Climate Change*, 140(1), 33-45. <https://doi.org/10.1007/s10584-013-0909-y>
- Anderson, C. M., DeFries, R. S., Litterman, R., Matson, P. A., Nepstad, D. C., Pacala, S., Schlesinger, W. H., Shaw, M. R., Smith, P., Weber, C., & Field, C. B. (2019). Field natural climate solutions are not enough. *Science*, 363(6430), 933-934. <http://dx.doi.org/10.1126/science.aaw2741>
- Balasoorya, H. N., Dassanayake, K. B., & Ajlouni, S. (2019). The impact of elevated CO<sub>2</sub> and high temperature on the nutritional quality of fruits - A short review. *American Journal of Agricultural Research*, 4(26), 1-9. <https://escipub.com/ajar-2018-12-1608/>
- Bhargava, S., & Mitra, S. (2021). Elevated atmospheric CO<sub>2</sub> and the future of crop plants. *Plant Breeding* 140, 1-11. <https://doi.org/10.1111/pbr.12871>
- Becker, C., & Kläring H. P. (2016). CO<sub>2</sub> enrichment can produce high red leaf lettuce yield while increasing most flavonoid glycoside and some caffeic acid derivative concentrations. *Food Chemistry*, 199, 736-745. <https://doi.org/10.1016/j.foodchem.2015.12.059>
- Bindi, M., Fibbi, L., & Miglietta, F. (2001). Free air CO<sub>2</sub> Enrichment (FACE) of grapevine (*Vitis vinifera* L.): II. Growth and quality of grape and wine in response to elevated CO<sub>2</sub> concentrations. *European Journal of Agronomy*, 14(2), 145-155. [https://doi.org/10.1016/S1161-0301\(00\)00093-9](https://doi.org/10.1016/S1161-0301(00)00093-9)
- Bisbis, M. B., Gruda, N., & Blanke, M. (2018). Potential impacts of climate change on vegetable production and product quality - a review. *Journal of Cleaner Production*, 170, 1602-1620. <https://doi.org/10.1016/j.jclepro.2017.09.224>
- Bradley, K. L., & Pregitzer, K. S. (2007). Ecosystem assembly and terrestrial carbon balance under elevated CO<sub>2</sub>. *Trends in Ecology and Evolution*, 22(10), 538-547. <https://doi.org/10.1016/j.tree.2007.08.005>
- Brito, F., Thaline, T., Pimenta, M., Henschel, J., Martins, S., Zsögön, A., & Ribeiro, D. (2020). Elevated CO<sub>2</sub> improves assimilation rate and growth of tomato plants under progressively higher soil salinity by decreasing abscisic acid and ethylene levels. *Environmental and Experimental Botany*, 176, 104050. <https://doi.org/10.1016/j.envexpbot.2020.104050>
- Brunori, E., Farina, R., & Biasi R. (2016). Sustainable viticulture: The carbon-sink function of the vineyard agro-ecosystem. *Agriculture, Ecosystems and Environment*, 223, 10-21. <https://doi.org/10.1016/j.agee.2016.02.012>
- Casierra-Posada, F. & Fischer, G. (2012). Poda de árboles frutales. In G. Fischer (Ed.), *Manual para el cultivo de frutales en el trópico* (pp. 169-185). Produmedios.
- Ceulemans, R., Janssens, I. A., & Jach, M. E. (1999). Effects of CO<sub>2</sub> enrichment on trees and forests: Lessons to be learned in view of future ecosystem studies. *Annals of Botany*, 84(5), 577-590. <https://doi.org/10.1006/anbo.1999.0945>
- Cortés, A. J., Restrepo-Montoya, M., & Bedoya-Canas, L. E. (2020). Modern strategies to assess and breed forest tree adaptation to changing climate. *Frontiers in Plant Science*, 11, 583323. <https://doi.org/10.3389/fpls.2020.583323>
- Cruz, J. L., Alves, A. A. C., LeCain, D. R., Ellis, D. D., & Morgan, J. A. (2016). Interactive effects between nitrogen fertilization and elevated CO<sub>2</sub> on growth and gas exchange of

- papaya seedlings. *Scientia Horticulturae*, 202, 32-40. <https://doi.org/10.1016/j.scienta.2016.02.010>
- DaMatta, F. M., Grandis, A., Arenque, B. C., & Buckeridge, M. S. (2010). Impacts of climate changes on crop physiology and food quality. *Food Research International*, 43, 1814-1823. <https://doi.org/10.1016/j.foodres.2009.11.001>
- De Zwart, H. F. (2012). Lessons learned from experiments with semi-closed greenhouses. *Acta Horticulturae*, 952, 583-588. <https://doi.org/10.17660/ActaHortic.2012.952.74>
- Dingkuhn, M., Luquet, D., Fabre, D., Muller, B., Yin, X., & Paul, M. J. (2020). The case for improving crop carbon sink strength or plasticity for a CO<sub>2</sub>-rich future. *Current Opinion in Plant Biology*, 56, 259-272. <https://doi.org/10.1016/j.pbi.2020.05.012>
- Dong, J., Gruda, N., Li, X., & Duan, Z. (2018). Effects of elevated CO<sub>2</sub> on nutritional quality of vegetables: A review. *Frontiers of Plant Science*, 9, 924. <https://doi.org/10.3389/fpls.2018.00924>
- Dong, J., Gruda, N., Li, X., Tang, Y., & Duan, Z. (2020). Impacts of elevated CO<sub>2</sub> on nitrogen uptake of cucumber plants and nitrogen cycling in a greenhouse soil. *Applied Soil Ecology*, 145, 103342. <https://doi.org/10.1016/j.apsoil.2019.08.004>
- Drake, B. G., & González-Meler, M. A. (1997). More efficient plants: a consequence of rising atmospheric CO<sub>2</sub>? *Annual Review of Plant Physiology and Plant Molecular Biology*, 48, 609-639. <https://doi.org/10.1146/annurev.arplant.48.1.609>
- Ebi, K. L., Anderson, C. L., Hess, J. J., Kim, S.-H., Loladze, I., Neumann, R. B., Singh, D., Ziska, L., & Wood, R. (2021). Nutritional quality of crops in a high CO<sub>2</sub> world: an agenda for research and technology development. *Environmental Research Letters*, 16, 064045. <https://doi.org/10.1088/1748-9326/abfcfa>
- Fischer, G., & Melgarejo, L. M. (2020). The ecophysiology of cape gooseberry (*Physalis peruviana* L.) - an Andean fruit crop. A review. *Revista Colombiana de Ciencias Hortícolas*, 14(1), 76-89. <https://doi.org/10.17584/rcch.2020v14i1.10893>
- Fischer, G., & Orduz-Rodríguez, J. (2012). Ecofisiología en frutales. In G. Fischer (Ed.), *Manual para el cultivo de frutales en el trópico* (pp. 54-72). Produmedios.
- Fischer, G., Ramírez, F., & Casierra-Posada, F. (2016). Ecophysiological aspects of fruit crops in the era of climate change. A review. *Agronomía Colombiana*, 34(2), 190-199. <https://doi.org/10.15446/agron.colomb.v34n2.56799>
- Gruda, N., Bisbis, M., & Tanny, J. (2019). Influence of climate change on protected cultivation: Impacts and sustainable adaptation strategies - A review. *Journal of Cleaner Production*, 225, 481e495. <https://doi.org/10.1016/j.jclepro.2019.03.210>
- Haokip, S. W., Shankar, K., & Lalringheta, J. (2020). Climate change and its impact on fruit crops. *Journal of Pharmacognosy and Phytochemistry*, 9(1), 435-438. <https://www.phytojournal.com/archives?year=2020&vol=9&issue=1&ArticleId=10464>
- Henson, R. (2011). *The rough guide to climate change* (3<sup>rd</sup> ed.). Penguin Books.
- Hiratsuka, S., Suzuki, M., Nishimura H., & Nada, K. (2015). Fruit photosynthesis in Satsuma mandarin. *Plant Science*, 241, 65-69. <https://doi.org/10.1016/j.plantsci.2015.09.026>
- Houston, L., Capalbo, S., Seavert, C., Dalton, M., Bryla, D., & Sagili, R. (2018). Specialty fruit production in the Pacific Northwest: adaptation strategies for a changing climate. *Climate Change*, 146, 159-171. <https://doi.org/10.1007/s10584-017-1951-y>
- IPCC. (2013). Climate change 2013: The physical science basis. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley

- (Eds.), *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- IPCC. (2019). *Summary for Policymakers*. In *Climate change and land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*. Intergovernmental Panel on Climate Change.
- Jackson, L. E., Wheeler, S. M., Hollander, A. D., O'Geen, A. T., Orlove, B. S., Six, J., Sumner, D. A., Santos-Martin, F., Kramer, J. B., Horwath, W. R., Howitt, R. E. T., & Tomich, T. P. (2011). Case study on potential agricultural responses to climate change in a California landscape. *Climate Change*, 109(Suppl 1), S407-S427. <https://doi.org/10.1007/s10584-011-0306-3>
- Jones, G. V., White, M. A., Cooper, O. R., & Storchmann, K. (2005). Climate change and global wine quality. *Climate Change*, 73, 319-343. <https://doi.org/10.1007/s10584-005-4704-2>
- Keutgen, N., Chen, A. I., & Lenz, F. (1997). Responses of strawberry leaf photosynthesis, chlorophyll fluorescence and macronutrient contents to elevated CO<sub>2</sub>. *Journal of Plant Physiology*, 150, 395-400. [https://doi.org/10.1016/S0176-1617\(97\)80088-0](https://doi.org/10.1016/S0176-1617(97)80088-0)
- Kimball, B. A., Idso, S. B., Johnson, S., & Rillig, M. C. (2007). Seventeen years of carbon dioxide enrichment of sour orange trees: final results. *Global Change Biology*, 13, 2171-2183. <https://doi.org/10.1111/j.1365-2486.2007.01430.x>
- Kizildeniz, T., Pascual, I., Irigoyen, J. J., & Morales, F. (2018). Using fruit-bearing cuttings of grapevine and temperature gradient greenhouses to evaluate effects of climate change (elevated CO<sub>2</sub> and temperature, and water deficit) on the cv. red and white Tempranillo. Yield and must quality in three consecutive growing seasons (2013-2015). *Agricultural Water Management*, 202, 299-310. <https://doi.org/10.1016/j.agwat.2017.12.001>
- Kläring, H. P., Hauschild, C., Heißner, A., & Bar-Yosef, B. (2007). Model-based control of CO<sub>2</sub> concentration in greenhouses at ambient levels increases cucumber yield. *Agricultural and Forest Meteorology*, 143, 208-216. <https://doi.org/10.1016/j.agrformet.2006.12.002>
- Kochhar, S. L., & Gujral, S. K. (2020). *Plant physiology: Theory and applications* (2<sup>nd</sup> ed.). Cambridge University Press. <https://doi.org/10.1017/9781108486392>
- Kumar, M., Sundaram, S., Gnansounou, E., Larroche, C., & Thakur, I. S. (2017). Carbon dioxide capture, storage and production of biofuel and biomaterials by bacteria: a review. *Bioresour Technol*, 247, 1059-1068. <https://doi.org/10.1016/j.biortech.2017.09.050>
- Larcher, W. (2003). *Physiological plant ecology*. Springer-Verlag. <https://doi.org/10.1007/978-3-662-05214-3>
- Leakey, A. D. B. (2009). Rising atmospheric carbon dioxide concentration and the future of C4 crops for food and fuel. *Proceedings of the Royal Society B*, 276, 2333-2343. <https://doi.org/10.1098/rspb.2008.1517>
- Leakey, A. D. B., Bishop, K. A., & Ainsworth, E. A. (2012). A multi-biome gap in understanding of crop and ecosystem responses to elevated CO<sub>2</sub>. *Current Opinion in Plant Biology*, 15, 228-236. <https://doi.org/10.1016/j.pbi.2012.01.009>
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., ..., & Zheng, B. (2018). Global Carbon Budget 2018. *Earth System Science Data*, 10, 2141-2194. <https://doi.org/10.5194/essd-10-2141-2018>



- Leung, D. Y., Caramanna G., & Maroto-Valer, M. M. (2014). An overview of current status of carbon dioxide capture and storage technologies. *Renewable and Sustainable Energy Review*, 39, 426-443. <https://doi.org/10.1016/j.rser.2014.07.093>
- Li, X., Zhao, J., Shang, M., Song, H., Zhang, J., Xu, X., Zheng, S., Hou, L., Li, M., & Xing, G. (2020). Physiological and molecular basis of promoting leaf growth in strawberry (*Fragaria × ananassa* Duch.) by CO<sub>2</sub> enrichment. *Biotechnology & Biotechnological Equipment*, 34(1), 905-917. <https://doi.org/10.1080/13102818.2020.1811766>
- López-Bellido, L. (2015). *Agricultura, cambio climático y secuestro de carbono*. Universidad de Córdoba.
- Luedeling, E., Girvetz, E. H., Semenov, M. A., & Brown, P. H. (2011). Climate change affects winter chill for temperate fruit and nut trees. *PLoS ONE*, 6(5), e20155. <https://doi.org/10.1371/journal.pone.0020155>
- Marín, M. del P., Andrade, H. J., & Sandoval, A. P. (2016). Fijación de carbono atmosférico en la biomasa total de sistemas de producción de cacao en el departamento del Tolima, Colombia. *Revista U.D.C.A Actualidad & Divulgación Científica*, 19(2), 351-360. <https://doi.org/10.31910/rudca.v19.n2.2016.89>
- Martínez-Lüscher, J., Morales, F., Sánchez-Díaz, M., Delrot, S., Aguirreolea, J., Gomès, E., & Pascual, I. (2015). Climate change conditions (elevated CO<sub>2</sub> and temperature) and UV-B radiation affect grapevine (*Vitis vinifera* cv. Tempranillo) leaf carbon assimilation, altering fruit ripening rates. *Plant Science*, 236, 168-176. <https://doi.org/10.1016/j.plantsci.2015.04.001>
- Menezes-Silva, P. E., Loram-Lourenço, L., Alves, R. D. F. B., Sousa, L. F., Almeida, S. E. D. S., & Farnese, F. S. (2019). Different ways to die in a changing world: Consequences of climate change for tree species performance and survival through an ecophysiological perspective. *Ecology and Evolution*, 2019, 1-21. <https://doi.org/10.1002/ece3.5663>
- Mishra, A. K., Agrawal, S. B., & Agrawal, M. (2019). Rising atmospheric carbon dioxide and plant responses: current and future consequences. In *Climate change and agricultural ecosystems* (pp. 265-306). Elsevier. <https://doi.org/10.1016/B978-0-12-816483-9.00011-6>
- Moretti, C. L., Mattos L. M., Calbo A. G., & Sargent S. A. (2010). Climate changes and potential impacts on postharvest quality of fruit and vegetable crops: A review. *Food Research International*, 43, 1824-1832. <https://doi.org/10.1016/j.foodres.2009.10.013>
- Morgan, J. A., LeCain, D. R., Pendall, E., Blumenthal, D. M., Kimball, B. A., Carrillo, Y., Williams, D. G., Heisler-White, J., Dijkstra, F. A., & West, M. (2011). C4 grasses prosper as carbon dioxide eliminates desiccation in warmed semi-arid grassland. *Nature*, 476, 202-206. <https://doi.org/10.1038/nature10274>
- Moriondo, M., Jones, G. V., Bois, B., Dibari, C., Ferrise, R., Trombi, G., & Bindi, M. (2013). Projected shifts of wine regions in response to climate change. *Climate Change*, 19(3-4), 825-839. <https://doi.org/10.1007/s10584-013-0739-y>
- Nobel, P. S. (1991). Achievable productivities of certain CAM plants: basis for high values compared with C3 and C4 plants. *New Phytology*, 119(2), 183-205. <https://doi.org/10.1111/j.1469-8137.1991.tb01022.x>
- Nobel, P. S. (1999). *Physicochemical & environmental plant physiology* (2<sup>nd</sup> ed.). Academic Press.
- Ouda, O. K. M., Raza. S. A., Nizami. A. S., Rehan. M., Al-Waked. R., & Korres. N. E. (2016). Waste to energy potential: a case study of Saudi Arabia. *Renewable and Sustainable Energy Reviews*, 61, 328-340. <https://doi.org/10.1016/j.rser.2016.04.005>

- Patil, P., & Kumar, K. (2017). Biological carbon sequestration through fruit crops (perennial crops - natural “sponges” for absorbing carbon dioxide from atmosphere). *Plant Archives*, 17(2), 1041-1046. [http://plantarchives.org/17-2/1041-1046%20\(3939\).pdf](http://plantarchives.org/17-2/1041-1046%20(3939).pdf)
- Paudel, I., Halpern, M., Wagner, Y., Raveh, E., Yermiyahu, U., Hoch, G., & Klein, T. (2018). Elevated CO<sub>2</sub> compensates for drought effects in lemon saplings via stomatal downregulation, increased soil moisture, and increased wood carbon storage. *Environmental and Experimental Botany*, 148, 117-127. <https://doi.org/10.1016/j.envexpbot.2018.01.004>
- Pérez, C., Nicklin, C., Dangles, O., Vanek, S., Sherwood, S., Halloy, S., Garrett, K., & Forbes, G. (2010). Climate change in the High Andes: Implications and adaptation strategies for small-scale farmers. *International Journal of Environmental, Cultural, Economic and Social Sustainability*, 6(5), 71-88. <https://doi.org/10.18848/1832-2077/CGP/v06i05/54835>
- Pérez-Jiménez, M., Hernández-Munuera, M., Piñero, M. C., López-Ortega, G., & del Amor F. M. (2017). CO<sub>2</sub> effects on the waterlogging response of ‘Gisela 5’ and ‘Gisela 6’ (*Prunus cerasus* × *Prunus canescens*) sweet cherry (*Prunus avium*) rootstocks. *Journal of Plant Physiology*, 213, 178-187. <https://doi.org/10.1016/j.jplph.2017.03.011>
- Polley, H. W. (2002). Implications of atmospheric and climate change for crop yield. *Crop Science*, 42, 131-140. <https://doi.org/10.2135/cropsci2002.1310>
- Prior, S. A., Runion, G. B., Torbert, H. A., Idso, S. B., & Kimball, B. A. (2012). Sour orange fine root distribution after seventeen years of atmospheric CO<sub>2</sub> enrichment. *Agricultural and Forest Meteorology*, 162-63, 85-90. <https://doi.org/10.1016/j.agrformet.2012.04.014>
- Pritchard, S. G., & Amthor, J. S. (2005). *Crops and environmental change*. Food Products Press, The Haworth Press.
- Rajan, R., Feza, M., Pandey, K., Aman, A., & Kumar, V. (2020). Climate change and resilience in fruit crops. In *Climate change and its effects on Agriculture* (pp. 337-354). Biotec Books.
- Ramírez, F., & Kallarackal, J. (2015). *Responses of fruit trees to global climate change*. Springer Briefs in Plant Science. Springer International Publishing. <https://doi.org/10.1007/978-3-319-14200-5>
- Reich, P. B., Hobbie, S. E., Lee, T., Ellsworth, D. S., West, J. B., Tilman, D., Knops, J. M. H., Naeem, S., & Trost, J. (2006). Nitrogen limitation constrains sustainability of ecosystem response to CO<sub>2</sub>. *Nature*, 440, 922-925. <https://doi.org/10.1038/nature04486>
- Sánchez, S., Morea, R., Serrano-Grijalva, L., Meco, A., & Sánchez-Andrés, R. A. (2015). Free Air CO<sub>2</sub> Enrichment (FACE) facility in a wetland to study the effects of elevated atmospheric carbon dioxide: System description and performance. *Wetlands*, 35, 193-205. <http://dx.doi.org/10.1007/s13157-014-0614-2>
- Sharma, S., Rana, V. S., Prasad, H., Lakra, J., & Sharma, U. (2021). Appraisal of carbon capture, storage, and utilization through fruit crops. *Frontiers in Environmental Science*, 9, 700768. <https://doi.org/10.3389/fenvs.2021.700768>
- Song, H., Li, Y., Xu, X., Zhang, J., Zheng, S., Hou, L., Xing, G., & Li, M. (2020). Analysis of genes related to chlorophyll metabolism under elevated CO<sub>2</sub> in cucumber (*Cucumis sativus* L.). *Scientia Horticulturae*, 261, 108988. <https://doi.org/10.1016/j.scienta.2019.108988>
- Stöckle, C. O., Marsal J., & Villar J. M. (2011). Impact of climate change on irrigated tree fruit production. *Acta Horticulturae*, 889, 41-52. <https://doi.org/10.17660/ActaHortic.2011.889.2>
- Sun, P., Mantri, N., Lou, H., Hu, Y., Sun, D., Zhu, Y., Dong, T., & Lu, H. (2012). Effects of elevated CO<sub>2</sub> and temperature on yield and fruit quality of strawberry (*Fragaria ananassa*

- Duch.) at two levels of nitrogen application. *PLoS ONE*, 7(79), e41000. <https://doi.org/10.1371/journal.pone.0041000>
- Taiz, L., Zeiger, E., Møller, I. M., & Murphy, A. (2017). *Fisiología e desenvolvimento vegetal* (6<sup>th</sup> ed.). Artmed.
- Tognetti, R., Raschi, A., Longobucco, A., Lanini, M., & Bindi, M. (2005). Hydraulic properties and water relations of *Vitis vinifera* L. exposed to elevated CO<sub>2</sub> concentrations in a free air CO<sub>2</sub> enrichment (FACE). *Phyton*, 45(3), 243-256. [https://www.zobodat.at/pdf/PHY\\_45\\_3\\_0243-0256.pdf](https://www.zobodat.at/pdf/PHY_45_3_0243-0256.pdf)
- Treutter, D. (2010). Managing phenol contents in crop plants by phytochemical farming and breeding-visions and constraints. *International Journal of Molecular Sciences*, 11, 807-857. <https://doi.org/10.3390/ijms11030807>
- Vélez, J. E., Polanía, W., & Beltrán, N. (2019). Efecto del régimen de riego en la producción de volátiles que incide en el aroma de la pera variedad Triunfo de Viena (*Pyrus communis* L.). *Revista Colombiana de Ciencias Hortícolas*, 13(3), 348-358. <https://doi.org/10.17584/rcch.2019v13i3.10920>
- Vélez-Sánchez, J. E., Balaguera-López, H. E., & Álvarez-Herrera, J. G. (2021). Effect of regulated deficit irrigation (RDI) on the production and quality of pear Triunfo de Viena variety under tropical conditions. *Scientia Horticulturae*, 278, 109880. <https://doi.org/10.1016/j.scienta.2020.109880>
- Wang, S. Y., Bunce J. A., & Maas, J. L. (2003). Elevated carbon dioxide increases contents of antioxidant compounds in field-grown strawberries. *Journal of Agriculture and Food Chemistry*, 51, 4315-4320. <https://doi.org/10.1021/jf021172d>
- Wei, Z., Du, T., Li, X., Fang, L., & Liu, F. (2018). Interactive effects of elevated CO<sub>2</sub> and N fertilization on yield and quality of tomato grown under reduced irrigation regimes. *Frontiers in Plant Science*, 9, 328. <https://doi.org/10.3389/fpls.2018.00328>
- Weiss, I., Mizrahi Y., & Raveh, E. (2010). Effect of elevated CO<sub>2</sub> on vegetative and reproductive growth characteristics of the CAM plants *Hylocereus undatus* and *Selenicereus megalanthus*. *Scientia Horticulturae*, 123, 531-536. <https://doi.org/10.1016/j.scienta.2009.11.002>
- Wohlfahrt, Y., Smith, J. P., Tittmann, S., Honermeier, B., & Stoll, M. (2018). Primary productivity and physiological responses of *Vitis vinifera* L. cvs. under Free Air Carbon dioxide Enrichment (FACE). *European Journal of Agronomy*, 101, 149-162. <https://doi.org/10.1016/j.eja.2018.09.005>
- Wu, T., Wang, Y., Yu1, C., Chiarawipa, R., Zhang, X., Han, Z., & Wu, L. (2012). Carbon sequestration by fruit trees - Chinese apple orchards as an example. *PLoS ONE*, 7(6), e38883. <https://doi.org/10.1371/journal.pone.0038883>
- Yohannes, H. (2016). A review on relationship between climate change and agriculture. *Journal of Earth Science and Climate Change*, 7(2), 335. <https://doi.org/10.4172/2157-7617.1000335>
- Zandalinas, S. I., Fritsch, F. B., & Mittler, R. (2021). Global warming, climate change, and environmental pollution: Recipe for a multifactorial stress combination disaster. *Trends in Plant Science*, 26(6), 588-599. <https://doi.org/10.1016/j.tplants.2021.02.011>